



Studies on Heterosis for Yield and its Component Traits in Brinjal (*Solanum melongena* L.)

**Anjana C. S. ^{a++*}, G. C. Yadav ^{a#}, Niharika Singh ^{a++},
Sandeep Kumar ^{a++} and Akhil Kumar Choudhary ^{a++}**

^a *Department of Vegetable Science, College of Horticulture and Forestry, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya-224229, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i113651

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109646>

Original Research Article

Received: 01/10/2023

Accepted: 09/12/2023

Published: 12/12/2023

ABSTRACT

A field trial was conducted to estimate the magnitude of heterosis in brinjal, for twenty-two characters, during *Rabi* 2021 at the Main Experiment Station, Department of Vegetable Science, Acharya Narendra Deva University of Agriculture and Technology, Ayodhya, Uttar Pradesh. 8 parental lines and the 28 F₁ hybrids obtained from the parental lines following half-diallel mating design, along with a standard variety, Pant Samrat were evaluated in randomized block design with three replications. Observations were recorded on twenty-two characters viz. days to 50% flowering, days to first fruit harvest, leaf area (cm²), plant height (cm), number of primary branches per plant length of fruit (cm), fruit diameter (cm), fruit length/circumference ratio, length of peduncle (cm), length of calyx (cm), number of fruits per cluster, number of fruits per plant, average fruit weight (g), fruit yield per plant (kg), crop duration, dry matter content (%), TSS (%), total phenol content (mg/100g), reducing sugars (%), non-reducing sugar (%), total sugars (%) and ascorbic

⁺⁺ P.G. Students;

[#] Associate Professor;

*Corresponding author: E-mail: anjanarohini97@gmail.com;

acid content (mg/100g). The study revealed that significant heterobeltiosis and standard heterosis were observed for all the characters under study. The crosses $P_2 \times P_4$ (42.42% and 25.12%) and $P_1 \times P_4$ (41.73% and 14.15%) exhibited considerable heterosis over the better parent and the standard variety respectively, which may be commercially exploited for cultivation under sodic soils due to their high yield and desirable fruit qualities.

Keywords: Brinjal (*Solanum melongena* L.); heterosis, better parent; standard variety; heterobeltiosis; standard heterosis.

1. INTRODUCTION

Brinjal (*Solanum melongena* L., $2n=2x=24$) is one of the most widely cultivated solanaceous vegetable crops in India. The fruit is botanically classified as a berry and the soft white fleshy placenta contains numerous small soft seeds embedded in it. The seeds are edible but are bitter as they contain nicotinoid alkaloids. Brinjal is referred as “vegetable of masses” as it is popular among people of all social strata (Patel et al. 2004). Due to the expanding population and rise in demand of the crop, there is need to further increase its productivity level. The crop improvement works in brinjal was started as early as 1900s and still exploration for a better cultivar is going on, as most of the commercial cultivars lack one or the other desirable traits [1]. Exploiting hybrid vigour has emerged as a promising approach for eggplant improvement. Nagai and Kida [2] were the first to notice brinjal hybrid vigour. Because of the cheap cost of F_1 seed production and the low seed demand per unit area, commercial exploitation of this phenomena has become achieved in the brinjal. With the growing popularity of F_1 hybrids in brinjal, it is critical to obtain such hybrids with outstanding quality and high yields.

Many scientists have reported on crop manifestations of heterotic impact for many economically relevant features. It is not possible to have a single cultivar that will suit all locales and tastes. There are differences in local preferences for characteristics such as color, size, form, and flavor and there are specific germplasm suited for specific region. Therefore, it is required to improve the locally preferred varieties with high yield potential and adaptation [3]. In India, brinjal has a huge amount of genetic diversity for agro-morphological characters [4]. This offers much scope for yield improvement through heterosis breeding. F_1 hybrids in India are more popular among farmers because of earliness, high yield potential, uniform maturity and high net return. In brinjal, only few breeding attempts have been undertaken for the development of potential hybrids as well as for

crop improvement by the exploitation of local germplasm in comparison to other Solanaceous crops [5]. The estimation of heterosis for fruit yield and quality parameters would be useful to judge the best hybrid combination for their commercial exploitation [6]. Keeping in view the above facts, the present study was undertaken with the objective to study the manifestation of heterosis in brinjal.

2. MATERIALS AND METHODS

The experimental materials for this study comprised of 37 genotypes which included 8 diverse parental lines, their 28 F_1 s, obtained by crossing in a half diallel fashion, and a standard variety Pant Samrat. The genotypes were evaluated in a randomized block design with three replications, following a spacing of 60 cm x 50 cm at the Main Experiment Station, Department of Vegetable Science, Acharya Narendra Deva University of Agriculture and Technology, Narendra Nagar, Ayodhya (Uttar Pradesh), Standard package of practices and management measures were followed to raise the crop. Data were recorded on five randomly selected plants in each treatment over replication for all the characters studied viz. days to 50% flowering, days to first fruit harvest, leaf area (cm^2), plant height (cm), number of primary branches per plant length of fruit (cm), fruit diameter (cm), fruit length/circumference ratio, length of peduncle (cm), length of calyx (cm), number of fruits per cluster, number of fruits per plant, average fruit weight (g), fruit yield per plant (kg), crop duration, dry matter content (%), TSS (%), total phenol content (mg/100g), reducing sugars (%), non-reducing sugar (%), total sugars (%) and ascorbic acid content (mg/100g). The magnitude of heterosis in hybrids was expressed as percentage of increase or decrease of a character over better parent and standard variety in the desirable direction and was calculated using the following formula as given by Hayes et al. [7].

$$\text{Heterobeltiosis (\%)} = \frac{\bar{F}_1 - \overline{BP}}{\overline{BP}} \times 100$$

$$\text{Standard heterosis} = \frac{\bar{F}_1 - \bar{SV}}{\bar{SV}} \times 100$$

Where,

\bar{F}_1 = mean value of F₁

\bar{BP} = mean value of better-parent and

\bar{SV} = mean value of standard variety.

3. RESULTS AND DISCUSSION

The range of heterosis for all the twenty-two characters studied are presented in Table 1. Negative heterosis is considered best for the maturity traits such as days to 50% flowering and days to first fruit harvest as it indicates earliness. For all the other characters studied, positive heterosis is beneficial. Heterosis for fruit yield per plant ranged from -33.67 % to 42.42 % over better parent and heterosis over standard variety varied from -51.89 to 25.12%. The result revealed that significant heterosis over better and standard varieties was observed for all the traits with minimum being in crop duration. However, significant desirable heterosis over both better parent and standard variety was not observed for crop duration and number of fruits per plant.

Table 2 shows the best three heterotic hybrids for each of the characters. It is observed that for the characters such as leaf area and length to circumference ratio better parent heterosis is not exhibited by any of the progenies. None of the hybrids showed both significant heterobeltiosis and standard heterosis for the traits such as crop duration and number of fruits per plant. The highest percentage of heterosis among the hybrids was observed over standard variety for the non-reducing sugar content. It implied that the hybrids P₆ x P₈ (317.86), P₃ x P₈ (291.67) and P₁ x P₂ (266.67) are more promising for the content of non-reducing sugar than the standard variety, Pant Samrat. Highest percentage of

heterosis over better parent was found for length of calyx (P₆ x P₇ (44.91), P₁ x P₃ (38.63), P₁ x P₄ (35.71)) and fruit yield per plant (P₂ x P₄ (42.42), P₁ x P₄ (41.73) and P₅ x P₇ (23.99)). Two crosses P₁ x P₄ and P₂ x P₄ were found significantly heterotic over better as well as standard variety for fruit yield per plant. The nature and magnitude of heterosis varied for different traits in various hybrid combinations. For maturity traits, negative heterosis is usually desirable, because this will promote earliness in hybrids compared to parents, thereby increasing the productivity per day per unit area. A close examination of heterosis values of the maturity trait *i.e.*, days to 50 per cent flowering and days to first fruit harvest (Table 2), revealed that thirteen and twelve F₁ hybrids respectively exhibited significant high heterosis in desirable direction over better parent. While, twenty-one and eleven crosses showed standard heterosis for days to 50% flowering and days to first fruit harvest, respectively over the standard variety. The significant heterosis for the maturity traits in desirable direction had been also reported by previous workers [8,9,10,11,12,13]. For all the other traits, positive values of heterosis is considered and the results are in proximity to the work done by Choudhary et al. [14], Bhatt et al. [15], Shitap et al. (2017) and Galani et al. [16]. In general, the hybrids with significant heterosis for yield also expressed significant heterosis either for fruit weight or for fruits per plant and this was in agreement with the findings by Singh et al. [17], Dutta et al. [18], Patel et al. [19], Dubey et al. [20] and Pandey et al. [21]. Crosses showing heterosis for other yield component did not necessarily show heterosis for fruit yield. This showed that heterosis depends upon nicking for genes. Similar results have also been reported by Reddy and Patel et al. [22] and Kumar et al. [23].

Table 1. Mean values of heterosis for the twenty-two characters studied

Characters	Range of Heterosis	
	Better parent	Standard variety
Days to 50% flowering	-32.33 to 1.64	-29.03 to 7.37
Days to first fruit harvest	-26.69 to 0.88	-20.54 to 15.82
Leaf area (cm ²)	-45.08 to 13.25	-27.30 to 78.12
Plant height (cm)	-36.04 to 14.75	-8.44 to 45.77
No: of primary branches per plant	-29.92 to 16.17	-15.80 to 35.48
Length of fruit (cm)	-37.95 to 37.31	-10.44 to 117.31
Circumference of fruit (cm)	-44.26 to 36.51	0.46 to 112.55
Length to circumference ratio	-57.89 to 11.96	-52.56 to 65.50
Length of pedicel (cm)	-37.05 to 33.85	-37.98 to 16.24
Length of calyx (cm)	-4.36 to 44.91	-15.18 to 47.11
No: of fruits per cluster	-14.59 to 22.44	-21.04 to 5.75

Characters	Range of Heterosis	
	Better parent	Standard variety
Average fruit weight	-17.83 to 36.04	-24.89 to 22.89
No: of fruits per plant	-39.44 to 7.01	-44.71 to 6.67
Fruit yield per plant (kg)	-33.67 to 42.42	-51.89 to 25.12
Crop duration	-5.16 to 4.32	-6.32 to 3.66
Dry matter content	-9.13 to 24.82	-14.30 to 24.55
TSS	-17.50 to 18.18	8.23 to 19.62
Total phenol content	-47.75 to 5.23	-46.06 to 5.25
Reducing sugars	-30.36 to 13.79	0.74 to 91.88
Non-reducing sugars	-69.83 to 18.98	5.95 to 317.86
Total sugars	-29.81 to 17.87	23.38 to 115.49
Ascorbic acid content	-34.37 to 22.31	-25.91 to 56.84

Table 2. Top three heterotic hybrids over better parent (BP) and standard variety (SV)

	Traits	BP	SV
1.	Days to 50% flowering	P ₆ X P ₇ (-32.33) P ₃ X P ₄ (-28.70) P ₃ X P ₇ (-27.35)	P ₁ X P ₂ (-29.03) P ₄ X P ₇ (-29.03) P ₅ X P ₇ (-29.03)
2	Days to first fruit harvest	P ₆ X P ₈ (-26.69) P ₃ X P ₈ (-20.20) P ₅ X P ₈ (-19.81)	P ₂ X P ₄ (-20.54) P ₁ X P ₄ (-19.53) P ₁ X P ₈ (-19.19)
3.	Leaf area	-	P ₁ X P ₃ (78.12) P ₂ X P ₃ (73.23) P ₃ X P ₆ (47.30)
4	Plant height	P ₇ X P ₈ (14.75) P ₆ X P ₈ (14.12) P ₅ X P ₇ (13.08)	P ₁ X P ₂ (45.77) P ₂ X P ₄ (45.27) P ₂ X P ₈ (45.18)
5	No: of primary branches per plant	P ₆ X P ₇ (16.17) -	P ₁ X P ₄ (35.48) P ₁ X P ₂ (32.44) P ₂ X P ₄ (14.82)
6	Length of fruit (cm)	P ₂ X P ₈ (37.31) P ₂ X P ₄ (35.20) P ₂ X P ₅ (19.50)	P ₂ X P ₈ (117.31) P ₂ X P ₄ (97.01) P ₅ X P ₈ (79.32)
7	Circumference of fruit (cm)	P ₄ X P ₈ (36.51) -	P ₆ X P ₇ (112.55) P ₁ X P ₇ (107.11) P ₅ X P ₇ (106.31)
8	Length to circumference ratio	-	P ₂ X P ₈ (65.50) P ₁ X P ₈ (60.92) P ₂ X P ₄ (56.60)
9	Length of pedicel (cm)	P ₅ X P ₆ (33.85) P ₃ X P ₆ (10.34) -	P ₃ X P ₆ (16.24) -
10	Length of calyx (cm)	P ₆ X P ₇ (44.91) P ₁ X P ₃ (38.63) P ₁ X P ₄ (35.71)	P ₂ X P ₃ (47.11) P ₃ X P ₄ (35.18) P ₁ X P ₃ (31.45)
11	No: of fruits per cluster	P ₃ X P ₆ (22.44) - -	-- - -
12	Average fruit weight	P ₂ X P ₄ (36.04) P ₁ X P ₄ (24.24) -	P ₆ X P ₇ (22.50) P ₃ X P ₇ (20.61) P ₂ X P ₇ (18.95)
13	No: of fruits per plant	- - -	- - -

	Traits	BP	SV
14	Fruit yield per plant (kg)	P ₂ X P ₄ (42.42) P ₁ X P ₄ (41.73) P ₅ X P ₇ (23.99)	P ₂ X P ₄ (25.12) - -
15	Crop duration	- - -	- - -
16	Dry matter content	P ₃ X P ₇ (24.82) P ₂ X P ₆ (22.65) P ₃ X P ₈ (21.17)	P ₂ X P ₆ (24.55) P ₂ X P ₅ (17.22) P ₃ X P ₇ (15.01)
17	TSS	P ₅ X P ₇ (18.18) P ₆ X P ₇ (13.75) P ₁ X P ₆ (13.13)	P ₄ X P ₈ (19.62) P ₁ X P ₈ (18.35) P ₄ X P ₇ (18.35)
18	Total phenol content	P ₂ X P ₃ (5.87) P ₃ X P ₇ (5.23) P ₃ X P ₈ (4.88)	P ₃ X P ₈ (5.25) P ₁ X P ₂ (4.77) P ₂ X P ₈ (4.65)
19	Reducing sugars	P ₂ X P ₃ (13.79) P ₂ X P ₄ (10.46) P ₁ X P ₆ (6.25)	P ₂ X P ₃ (91.88) P ₄ X P ₅ (62.73) P ₂ X P ₅ (60.89)
20	Non-reducing sugar	P ₆ X P ₈ (18.98) P ₆ X P ₇ (18.56) P ₃ X P ₈ (11.53)	P ₆ X P ₈ (317.86) P ₃ X P ₈ (291.67) P ₁ X P ₂ (266.67)
21	Total sugar	P ₆ X P ₈ (17.87) P ₃ X P ₈ (9.40) -	P ₆ X P ₈ (115.49) P ₁ X P ₄ (103.94) P ₃ X P ₈ (100.00)
22	Ascorbic acid content	P ₂ X P ₄ (22.31) - -	P ₄ X P ₈ (56.84) P ₃ X P ₈ (56.78) P ₂ X P ₈ (55.28)

4. CONCLUSION

The above findings indicated that some pureline has strong heterotic capability compared to other ones during hybridization process. This may be due to diverse parent and favorable cross combination. As the performance of hybrids developed upon the heterotic capability of the parents involved, from economic point of view it will be useful to select and utilize the pure line with strong heterotic capability for important traits associated with yield in order to achieve higher fruit yield in F₁ hybrids through exploitation of heterosis. Since earliness, desirable fruit shape and color are the important considerations for choice of elite high yielding F₁ hybrids, the decision for final selection of a hybrid for commercial cultivation should also take into account the earlier three factors *i.e* earliness, fruit shape and colour traits along with the latter *i.e* high fruit yield and other desirable qualitative attributes for market preference.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Prabakaran S. Evaluation of local types of egg plant (*Solanum melongena* L.). Thesis. M.Sc. (Hort.), Agricultural College and Research Institute, TNAU, Madurai; 2010.
2. Nagai K, Kida S. Experiments on hybridization of various strains of *Solanum melongena*. Japan. J. Genet. 1926;4:10-30.
3. Akpan N, Ogbonna P, Onyia V, Okechukwu E, Atugwuand A, Dominic IO. Studies on the variability and combining ability for improved growth and yield of local eggplant genotypes (*Solanum melongena* L.). Not. Sci. Bio. 2016;226-231.
4. Sidhu AS, Bal SS, Behera TK, Rani M. An outlook in hybrid eggplant breeding. J. New Seeds. 2005;6(2-3):15-29.
5. Barik S, Ponnam N, Acharya G, Singh TH, Kumari M. Genetics of growth and yield attributing traits of brinjal (*Solanum melongena* L.) through six generation mean analysis. J. Hortic. Sci. 2022;17(1): 41-50.
6. Prabhu M, Natarajan S, Pugalendhi L. Studies on heterosis and mean

- performance in brinjal (*Solanum melongena* L.). Veg. Sci. 2005;32(1):86-87.
7. Hayes HK, Immer FR, Smith DC. Methods of plant breeding. McGraw Hill Co. Inc. 1955;2:52-65.
 8. Kumar, A, Sharma V, Jain BT, Kaushik P. Heterosis breeding in eggplant (*Solanum melongena* L.): Gains and provocations. Plants. 2020;9(3):403.
 9. Deshmukh SB, Narkhede GW, Gabale LK, Dod VN. Hybrid vigour in brinjal (*Solanum melongena* L.). The Bioscan. 2015;10(2): 869-876.
 10. Mistry CR, Kathiria KB, Sabolu S, Kumar S. Heterosis and inbreeding depression for fruit yield attributing traits in eggplant. Current Plant Biology. 2018;16:27-31.
 11. Dishri M, Mishra HN, Senapati N, Talekar NS, Sahu GS. Estimation of heterosis for earliness, yield and its components in brinjal (*Solanum melongena*). Plant Arch., 2018;18(2):1405-1410.
 12. Sharma TK, Pant SC, Kumar S, Paliwal A, Bahuguna P, Badhani HC. Combining ability studies in brinjal (*Solanum melongena* L.). Int. J. Bio-res. Stress Manag. 2016;7(6):1225-1231.
 13. Rani M, Kumar S, Kumar M. Estimation of heterosis for yield and its contributing traits in brinjal. J. Environ. Biol. 2018;39(5):710-718.
 14. Chaudhari BN, Patel AI, Vashi JM. Study on Heterosis over Environments in Brinjal (*Solanum melongena* L.). Int. J. Curr. Microbiol. App. Sci. 2020;9(7):3358-3367.
 15. Bhatt PK, Zapadiya VJ, Sapovadiya MH, CP C. Estimation of heterobeltiosis and standard heterosis for fruit yield and its attributes in brinjal (*Solanum melongena* L.). J. Pharmacogn. Phytochem. 2019;8(4): 1384-1388.
 16. Galani SN, Senjaliya HJ, Mungra KS, Gorfad PS. Heterosis for fruit yield and its component traits in brinjal (*Solanum melongena* L.). Trend. Biosci. 2015;8(11): 2952-2956.
 17. Singh S. Study of heterosis for vegetative and quantitative traits in brinjal (*Solanum melongena* L.). The Pharma Journal. 2021; 10(12):335-338.
 18. Dutta T, Banerjee S, Bhattacharjee T, Maurya PK, Dutta S, Chattopadhyay A, Hazra P. Heterosis breeding for improving quality traits in brinjal for export in the tropics. Indian J. Horticulture. 2019;76(3): 438-445.
 19. Patel AA, Gohil DP, Dhruve JJ, Damor HI. Heterosis for fruit yield and its quality characters in brinjal (*Solanum melongena* L.). J. Pharmacogn. Phytochem. 2017; 6(6):975-978.
 20. Dubey R, Das A, Ojha MD, Saha B, Ranjan A, Singh PK. Heterosis and combining ability studies for yield and yield attributing traits in brinjal (*Solanum melongena* L.). The Bioscan. 2014;9(2): 889-894.
 21. Pandey S, Mishra S, Kumar N, Yadav GC, Pandey VP. Studies on heterosis, combining ability and gene advance for the quantitative characters in brinjal or egg plant (*Solanum melongena* L.). J. Pharmacogn. Phytochem. 2019;8(1), 19-22.
 22. Reddy EEP, Patel AI. Heterosis studies for yield and yield attributing characters in brinjal (*Solanum Melongena* L.). J. Recent Adv. Agri. 2014;2(2):175-180.
 23. Kumar SR, Arumugam T. Gene action and combining ability analysis in brinjal (*Solanum melongena* L.). J. Hort. Sci. 2013;8(2):249-254.

© 2023 Anjana et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/109646>