



Impact of Nutrient Management Practices on Exchangeable K, Ca, Mg, CEC and Correlations with Soil Properties Under Maize-wheat Cropping System

Madhuri Toppo ^a, Asha Kumari Sinha ^a,
Dhirendra Kumar Shahi ^a and Ajay Kumar Upadhyay ^{b*}

^a Department of Soil Science & Agricultural Chemistry, Birsa Agricultural University, Ranchi-834006 (Jharkhand), India.

^b Department of Soil Science, Dr. Rajendra Prasad Central Agricultural University, Pusa, Samastipur-848125 (Bihar), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i7964>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116989>

Original Research Article

Received: 22/03/2024

Accepted: 27/05/2024

Published: 03/06/2024

ABSTRACT

In order to study the impact of nutrient management practices on exchangeable K, Ca, Mg, CEC and its correlations with soil properties in two depths (0-15 cm and 16-30 cm) under maize-wheat cropping system, a long term experiment was conducted at farm research centre of department of

*Corresponding author: E-mail: ajpusasoil.339@gmail.com;

Cite as: Toppo, M., Sinha, A. K., Shahi, D. K., & Upadhyay, A. K. (2024). Impact of Nutrient Management Practices on Exchangeable K, Ca, Mg, CEC and Correlations with Soil Properties Under Maize-wheat Cropping System. *Journal of Advances in Biology & Biotechnology*, 27(7), 38–50. <https://doi.org/10.9734/jabb/2024/v27i7964>

soil science falling under the jurisdiction of Birsa Agricultural University. The experiment consisted of nine treatments replicated thrice in a randomized block design were selected for the study. The selected treatments were T₁-control, T₂-100% N only where 100% N was applied through Urea fertilizers, T₃- FYM where 100% N was supplied through FYM (full dose 22 t/ha), T₄-100% NP here 110 and 90 kg ha⁻¹ were supplied through Urea and SSP fertilizers, respectively, T₅-100% NPK here 110:90:70 kg ha⁻¹ were supplied through Urea: SSP: MOP fertilizers, T₆- ½ (N+FYM)+P (A-X/2)+K (B-Y/2) (INM) where 50% of the nutrient need was supplied through Urea, SSP & MOP fertilizer and the rest 50% need through FYM and T₇-100% NPK + Lime where NPK applied through Urea, SSP & MOP fertilizer along with lime application in LR once in 4 years, T₈-Lime+ FYM+ P (A-X)+K (B-Y), the 100% N need of the crop was met by application of FYM whereas the P & K need was met by the application of SSP & MOP fertilizer, T₉-Lime+N the 100% N dose was supplied through Urea along with lime. The results indicated that exchangeable K was found to be significantly highest in T₈ than rest of the treatments followed by T₇ and T₅ while lowest was in T₄ treatment in both depths respectively. The exchangeable Ca²⁺ was found to be highest in treatment T₇ followed by T₈ and T₃ than rest of the other treated plot while lowest amount was found in T₂ in both the depths. The trend of exchangeable magnesium was found to be significantly different in 0-15 cm and 15-30 cm soil depths as the highest amount was observed in treatments such as T₈ followed by T₃ and T₆ in surface depths (0-15 cm) while in sub-surface depths (15-30 cm) the highest amount was found in treatments T₃ followed by T₈ and T₆ while the lowest amount was found in T₂ treated plot in both depths. Cation exchangeable capacity (CEC) is one the vital soil parameters concerning to soil fertility measured in terms of c mol (p+) kg⁻¹ of soil and it is significantly varied in surface soils as well as in sub-surface soil depths. The CEC was significantly higher in FYM treated plot as compared to Non-FYM treated plot. Therefore, the highest CEC was recorded in T₃ treated plot than rest of the other treatment while the lowest one was recorded in T₂ plot in surface soils as well as sub-surface soils. The correlation coefficient (r) values worked out among the various soil parameters at these depth of soil where pH was significantly and positively correlated with exchangeable Ca²⁺ and exchangeable Mg²⁺. Similarly organic carbon shows positive correlation with exchangeable Mg²⁺ and exchangeable K⁺. Whereas EC and CEC shows non-significant but positive correlation with the parameters taken under study. The result of the present study will be highly useful for improving the soil fertility by using suitable and judicious combination of nutrient management with various cropping system side by side and hence provides a scientific scope for further research work.

Keywords: Nutrient management; soil properties; synthetic fertilizers; Indian agriculture.

1. INTRODUCTION

“India and its sub-continent has made tremendous progress in agriculture after the green revolution or after the 1960’s. The success of agricultural production has been widely depend on the use of several inputs such as high yielding varieties, synthetic fertilizers, irrigation etc. Among these inputs, use of fertilizers arise as a vital input and become the kingpin for the transformation of indian agriculture from subsistence to surplus” (Ravindra Chary et al., 2019). “After the resounding success of the green revolution, a decline in the growth rate of food production is seen during the recent past in respect of crop productivity and input use response. Presently, the major concern in agriculture is to arrest any further decline in crop productivity and soil quality. Although high yielding nutrient responsive crop varieties resulted in higher productivity, it led to the

overexploitation of soil reserves and other resources. In general, Indian soils are poor in fertility, as these have been consistently depleted of their finite nutrient resources due to continuous cultivation for centuries, adoption of modern agricultural technologies, and imbalanced use of fertilizers and poor use efficiency of fertilizers” [1,2]. “It is essential to enhance the productivity of prominent crops of the country like paddy, wheat, maize etc. through location- specific nutrient management practices. To augment major food crops production, Food and Agriculture Organization (FAO) conceptualized the idea of plant nutrients in a crop and cropping system for better resource use. It is not only a reliable way of obtaining fairly higher yields with substantial fertilizer economy, but also a concept that is ecologically sound leading to sustainable agriculture. None of the nutrient sources alone can meet the total plant nutrients. Integration of inorganic and organic

sources nutrient sources is the most efficient way to supply plant nutrients for sustained crop productivity and improved of soil fertility” [3]. Generally cereal crops like maize, wheat, paddy etc. belongs to graminaceae being, exhaustive crops, capable to removes large amount of nutrients such as N, P, exchangeable K⁺, Ca²⁺, Mg²⁺ from soil.

“It is therefore necessary to judiciously manage the inflow of organic sources of nutrients, and their integration with fertilizers, and organic manure. Application of organic materials along with inorganic fertilizers leads to increased productivity of the system and sustained soil health for a longer period” [4]. Due to escalation of fertilizer prices and associated environment problem the crisis has necessitated in search for alternative sources of manures for integrated nutrient management, which includes inorganic, organic manures, lime etc. to sustain the cereal based cropping system.

2. MATERIALS AND METHODS

2.1 Experimental Details

The field study was as a part of an ongoing Permanent Manurial Trial (PMT) initiated in 1956 on the maize-wheat cropping system on acidic red loamy soil at the Experimental Farm (85° 19' E longitude, 23° 17' N latitude) of Birsa Agricultural University, Kanke, Ranchi, Jharkhand. The experiment was laid out in

Randomized Block Design with 3 replications and 9 treatments in each replication having a plot size of 10 m². The treatment details are presented in Table 1. The recommended dose of N: P₂O₅: K₂O are 110, 90, and 70 kg ha⁻¹, respectively, since 1976. Lime is applied as per lime requirement in treatments once in four years. FYM is applied based on N content as per the treatments @ 22 t ha⁻¹ 15 days before the sowing of each crop. Phosphorous and potassium are applied as basal and nitrogen is applied in splits for both the crops. The sources of N, P, and K were urea, single superphosphate (SSP), and muriate of potash (MOP), respectively. The varieties Suwan composite and HD 2967 for maize and wheat were used as a test crop respectively.

2.2 Collection and Analysis of Soils

A total of 54 composite soil samples were collected at two different depths viz. 0-15 cm (surface depth) & 16-30 cm (sub-surface depth) after the harvest of crop from each plot of the experimental field of the nine selected treatments replication wise. The soil samples collected were air-dried in shade, gently ground using wooden pestle and mortar to pass through 2 mm sieve. The sieved samples were preserved in stoppered plastic containers for further analysis. Processed and well-dried soil samples were used to analyze the various forms of potassium by using the standard soil:extractant ratio with standard and specific methods for each.

Table 1. Details of the various treatments of the long-term field experiment

Treatment No.	Treatment description	Particulars
T ₁	Control	No fertilizer, manure, or lime
T ₂	100 % N	110 kg N ha ⁻¹ as urea
T ₃	FYM	FYM was applied @ 22 t ha ⁻¹ , 15 days before sowing of Maize-wheat crop
T ₄	100 % NP	110 kg N ha ⁻¹ as Urea + 90 kg P ₂ O ₅ kg ha ⁻¹ as Single Super Phosphate (SSP)
T ₅	100 % NPK	110 kg N ha ⁻¹ as urea + 90 Kg P ₂ O ₅ Kg ha ⁻¹ as SSP+70 Kg K ₂ O as MOP
T ₆	½ (N+FYM) + P _(A-X/2) + K _(B-Y/2) (INM)	50% N substituted through FYM and NPK (55.0 kg N ha ⁻¹ as urea + 55.6 kg P ₂ O ₅ kg ha ⁻¹ as SSP+42.9 kg K ₂ O as MOP
T ₇	100 % NPK + Lime	Lime as per LR (once in four years) + NPK (110 kg N ha ⁻¹ as urea + 90 kg P ₂ O ₅ kg ha ⁻¹ as SSP+70 Kg K ₂ O as MOP)
T ₈	Lime+ FYM+P _(A-X) +K _(B-Y)	Lime as per LR (once in four years) + FYM (22 t ha ⁻¹) + 55.6 kg P ₂ O ₅ kg ha ⁻¹ as SSP+42.9 kg K ₂ O as MOP
T ₉	Lime + N	Lime as per LR (once in four years) + N @110 kg N ha ⁻¹ as Urea

2.3 Methods Used for Analysis of Soil Parameters

Table 2. Methods Used for Soil Parameters Analysis

Sl.No	Soil Parameters	Methods
1.	Exchangeable Ca	Versanate Titration
2.	Exchangeable Mg	Versanate Titration
3.	Cation Exchange Capacity (CEC)	1N Ammonium Acetate
4.	pH	Soil:Water Suspension (1:2)
5.	Electrical Conductivity (EC)	Soil:Water Suspension (1:2)
6.	Organic Carbon (OC)	Walkley and Black
7.	Water Soluble K	Flame Photometer
8.	Available K	Flame Photometer
9.	Exchangeable K	Available K-Water Soluble K

2.4 Detailing of Methods for Soil Exchangeable Calcium, Magnesium and Exchangeable Potassium

Exchangeable calcium and magnesium: Exchangeable Ca plus Mg were extracted by using neutral normal ammonium acetate and estimated by complexometric titration method (versanate titration method) as described in Baruah and Barthakur [5]. The concentrations of calcium and magnesium ions were determined by complexometric titration with EDTA (Hesse, 1971), using calcon red indicator for calcium and erichrome black T (EBT) indicator for calcium plus magnesium. Magnesium concentration was estimated by subtraction.

Water soluble K: Water soluble K was determined by flame photometer after extraction of soil with doubled distilled water (Hanway and Heidal, 1952) in soil solution ratio 1:5 (w/v).

Available K: Available potassium was determined by Flame photometer after extraction of soil with neutral ammonium acetate (Jackson 1973) in soil to solution ratio 1:5 (w/v).

Exchangeable K: Exchangeable K was estimated by deducting the value of water soluble K from available K.

2.5 Statistical Analysis

The data sets were processed for analysis of variance as applicable to randomized block design using the least significant difference as described in Gomez and Gomez (1984). Treatment means were compared at 1% or 5% level of significance. Correlation studies were carried out among the parameters of soil using Microsoft Excel.

3. RESULTS AND DISCUSSION

The present investigation entitled "Impact of Nutrient Management Practices on Exchangeable K, Ca, Mg, CEC and Its Correlations with Soil Properties Under Maize-Wheat Cropping System" was conducted during 2017-18 on the Permanent Manurial Trial at Experimental farm of Birsa Agricultural University, Ranchi. Attempts have been made to present different observations on changes on soil exchangeable potassium, calcium, magnesium fraction as well as CEC and their correlations with soil properties at two depths i.e. surface soil (0-15 cm) and sub-surface (16-30 cm) have been discussed. Simultaneously, assessment of water soluble and available potassium also carried out in order to assess the exchangeable potassium.

3.1 Impact of Nutrient Management Practices on Different forms of Potassium

To assess the exchangeable potassium it is necessary to assess the water soluble and available potassium from the surface soil and sub-surface depth of soil which is presented in tables and graphs.

3.1.1 Water soluble potassium

Perusal of data (Table 3 & Table 4) indicated that under different nutrient- management treatments, the concentration of water soluble K in the surface soil (0-15 cm depth) ranged from 6.4 to 24.24 kg ha⁻¹ and sub-surface soil (16-30 cm) ranged from 6.03 to 23.91 kg ha⁻¹. The highest amount of water soluble was observed in treatment T₈ which is at par with T₅ and T₇ and followed by T₂. The lowest amount was observed in treatment T₄. Almost similar trend was found in both depths w.r.t. various treatment combination

except T₂ and T₆. On critical examination of data it was also observed that water soluble K was numerically lower in sub-surface (16-30 cm) soil than surface (0-15 cm) soil. This could be attributed to release of labile K from organic residues, application of K containing fertilizers and upward translocation of K from lower soil depths with capillary rise of groundwater Lakaria et al. [6] and Mazumdar et al. [7]. This may be due to continuous cropping without addition of K for the last 62 years. Similar findings were reported by [8]. It was also observed that water soluble K of 100% NP treated plot was also significantly lower than that of 100% N treated plot. This may be due to the yield of 100% NP treated plot was more than that of 100% N treated plot. The higher content of water soluble K was recorded in treatment-8 where fertilizer with FYM was applied. This may be due to, in this treatment, K was applied in higher quantity (in the form of 100% K₂O+ FYM) than others since the inception of this experiment. The increased status of water soluble K with Lime, FYM, P₂O₅, K₂O treatment may also be due to stimulating effect of FYM in reducing K fixation, thereby bringing in more K into available form [9] and similar findings reported by Kumari [10].

3.1.2 Available potassium

Perusal of data (Table 3 & Table 4) indicated that under different nutrient- management treatments, the concentration of Available K in the surface soil ranged from 62.9 to 225 kg ha⁻¹ and in sub-surface soil ranged from 57.7 to 219.8 kg ha⁻¹. In surface soil, the highest Available K status was found when plot was treated with full dose of Lime, FYM, P₂O₅ and K₂O fertilizers (T₈), which was significant higher than other treatments. It was followed by 100% NPK (T₅) treated plot. It was also observed that Available K of 100% NP treated plot (T₄) was significantly lower than all the treatments. Similar trend of Available K status was also found in sub-surface soil. The higher content of water soluble K was recorded in treatment-8 where fertilizer with FYM was applied. This may be due to, in this treatment, K was applied in higher quantity (in the form of 100% K₂O+ FYM) than others since the inception of this experiment. The increased status of water soluble K with Lime, FYM, P₂O₅, K₂O treatment may also be due to stimulating effect of FYM in reducing K fixation, thereby bringing in more K into available form [9] and similar findings reported by Kumari [10]. While the lower availability in T₄ was due to continuous cropping

without addition of K. Similar findings were reported by [8]. On critical examination of data it was also observed that water soluble K was numerically lower in sub-surface (16-30 cm) soil than surface (0-15 cm) soil. This could be attributed to release of labile K from organic residues, application of K containing fertilizers and upward translocation of K from lower soil depths with capillary rise of groundwater Lakaria et al. [6] and Mazumdar et al. [7].

3.1.3 Exchangeable potassium

Perusal of data (Table 3 & Table 4) indicated that under different nutrient- management treatments, the concentration of Exchangeable K in the surface soil ranged from 56.5 to 200.76 kg ha⁻¹ and in the sub-surface soil ranged from 51.67 to 195.89 kg ha⁻¹. In surface soil, the highest Exchangeable K was found when plot was treated with full dose of Lime, FYM, P₂O₅ and K₂O fertilizers (T₈), which was significant higher than other treatments. It was followed by 100% NPK (T₅) treated plot. It was also observed that Exchangeable K of 100% NP treated plot (T₄) was significantly lower than all the treatments. Similar trend of Exchangeable K status was also found in sub-surface soil. The highest exchangeable K concentration under Lime+FYM+P₂O₅+K₂O(T₈) was also might be due to the fact that addition of FYM could increase the CEC of soil, which can hold more exchangeable K and convert K from Non-exchangeable form to exchangeable form, consequent to mass action effect. Bhattacharyya et al. [11] reported that higher amount of K attribute to the process of structural K release through increasing the area of exchangeable surfaces, and due to the acceleration weathering of the interlayer K by application of FYM. The higher concentration of Exchangeable K under K fertilized plots in surface soil could be attributed to the addition of K through plant residues, manure and fertilizers [12] (Jadhao et al., 2018) in the plot. While the lower availability in T₄ was due to continuous cropping without addition of K. Similar findings were reported by [8]. It was also observed that the concentration of exchangeable K was numerically lower in sub-surface soil as compare to surface soil in all the treatments which may be due to comparatively more weathering vegetation and supply of K from organic residues in surface layer in lower depth. Sharma et al. [13] and Mazumdar et al. [7]. Similar findings was reported by Sheela Swamy (2006), Lakaria et al. [6] and Divya et al., (2016).

Table 3. Effect of nutrient management practices on different forms of K in surface soil

		Surface soil 0-15 cm		
Treatments		Water soluble K (kg ha ⁻¹)	Exchangeable K (kg ha ⁻¹)	Available K (kg ha ⁻¹)
T ₁	Control	9.14	79.56	88.7
T ₂	100 % N	13.8	111.3	125.1
T ₃	FYM	11.5	98.23	109.7
T ₄	100 % NP	6.40	56.5	62.9
T ₅	100 % NPK	14.89	116.11	131
T ₆	½ (N+FYM) + P (A-X/2) + K (B-Y/2) (INM)	12.63	103.07	115.7
T ₇	100 % NPK + Lime	20.90	83.6	104.5
T ₈	Lime +FYM+P(A-X)+K(B-Y)	24.24	200.76	225
T ₉	Lime + N	8.23	67.27	75.5
CD(P=0.05)		0.90	3.85	4.04
CV(%)		4.37	2.23	2.09

Table 4. Effect of nutrient management practices on different forms of K in sub-surface soil

		Subsurface soil 16-30 cm		
Treatments		Water soluble K(kg ha ⁻¹)	Exchangeable K (kg ha ⁻¹)	Available K (kg ha ⁻¹)
T ₁	Control	8.67	75.53	84.2
T ₂	100 % N	12.85	106.85	119.7
T ₃	FYM	11	93.47	104.6
T ₄	100 % NP	6.03	51.67	57.7
T ₅	100 % NPK	13.3	113	126.3
T ₆	½ (N+FYM) + P (A-X/2) + K (B-Y/2) (INM)	12.11	98.39	110.5
T ₇	100 % NPK + Lime	20.45	76.95	97.4
T ₈	Lime +FYM+P(A-X)+K(B-Y)	23.91	195.89	219.8
T ₉	Lime + N	8.10	63.2	71.3
CD(P=0.05)		0.97	5.60	5.72

3.2 Impact of Nutrient Management Practices on Exchangeable Calcium, Magnesium and Cation Exchange Capacity

3.2.1 Exchangeable calcium

Under different nutrient- management treatments, the concentration of Exchangeable calcium in the surface soil ranged from 2.5 to 8.1 meq 100g⁻¹ and in sub-surface soil ranged from 2.1 to 7.7 meq 100g⁻¹ respectively. In the surface soil, the amount of exchangeable calcium was 8.1 meq 100g⁻¹ (T₇) when plot was treated with 100 % NPK, Lime fertilizer which was significant higher than other plots. It was followed by (T₈) Lime, FYM, P₂O₅, K₂O treated plot. It was also observed that Exchangeable calcium of 100% N treated plot (T₂) was significantly lower (2.5 meq 100g⁻¹) than all the treatments. The exchangeable Ca status of continuously fertilized acid soil increased due to the application of lime

and FYM. Application of lime increased exchangeable Ca whereas FYM application increased Prasad et al. [14] suggested that the high level of Ca in FYM treated plots was due to increased biomass production and its incorporation in the soil. Babu et al. [15] also reported increase in exchangeable Ca with the application of FYM. An increase in pH of the soil is also associated with the increased availability of Ca and Mg ions [16]. Similar findings reported by Verma [17]. While the lower amount of exchangeable calcium might be due reduced pH as well as continuous removal of Ca by the crops. Several authors have reported that in multiple cropping systems, higher fertilizer levels decreased the availability of calcium in soils. On critical examination of data it was also observed that exchangeable Ca was decreased in sub-surface (16-30 cm) soil than surface (0-15 cm) soil and exchangeable Ca of surface and sub-surface soil followed the same trend.

3.2.2 Exchangeable magnesium

Under different nutrient-management treatments, the concentration of exchangeable Magnesium in the surface soil ranged from 0.7 to 3.9 meq 100g⁻¹ and in the sub-surface soil ranged from 0.13 to 3.60 meq 100g⁻¹. In the surface soil, the amount of exchangeable magnesium was 3.9 meq 100g⁻¹ (T₈) Lime, FYM, P₂O₅, K₂O treated plot which was significantly higher than other plots. It was followed by (T₃) FYM treated plot. It was also observed that exchangeable magnesium of 100% N treated plot (T₂) was significantly lower (0.7 meq 100g⁻¹) than all the treatments. Exchangeable Mg content were lowest for N treatment, which might be due reduced pH as well as continuous removal of Mg by the crops. Several authors have reported that in multiple cropping systems, higher fertilizer levels decreased the availability of Mg in soils. The exchangeable Mg status of continuously fertilized acid soil increased due to the application of lime and FYM. Application of lime increased exchangeable Mg whereas FYM application increased Prasad et al. [18] suggested that the high level of Mg in FYM treated plots was due to increased biomass production and its incorporation in the soil. Babu et al. [15] also reported increase in exchangeable Mg with the application of FYM. An increase in pH of the soil is also associated with the increased availability of Mg ions [16]. Whereas in sub-surface soil, the amount of exchangeable magnesium was 3.60 meq 100g⁻¹ (T₃) FYM, treated plot which was significantly higher than other plots. It was followed by (T₈) Lime, FYM, P₂O₅, K₂O treated plot. It was also observed that exchangeable magnesium of 100% N treated plot (T₂) was significantly lower (0.13 meq 100g⁻¹) than all the treatments. On critical examination of data it was

also observed that exchangeable Mg was decreased in sub-surface (16-30 cm) soil than surface (0-15 cm) soil.

3.2.3 Cation exchange capacity

Under different nutrient- management treatments, the concentration of CEC in the surface soil ranged from 6.7 to 13.2 cmol (p+) kg⁻¹ and in sub-surface soil ranged from 5.83 to 11.6 cmol (p+) kg⁻¹ respectively. The highest CEC in surface soil was 13.2 cmol (p+) kg⁻¹ of soil recorded in (T₃), when plot was treated with FYM fertilizer which was significant higher than other plots. It was followed by (T₈) treated plot 12.8 cmol (p+) kg⁻¹ of soil. It was also observed that CEC of 100% N treated plot (T₂) was significantly lower (6.7meq 100g⁻¹) than all the treatments. The CEC value of soils was comparatively high in the FYM-treated plots, which might be due to the higher organic colloids in these plots. Moreover, the increase in CEC with the addition of FYM can be attributed to the increase in root biomass and crop residues production and their incorporation in the soil. Similar findings were reported by Prasad et al. [14] and Sharma (2004) and similar findings reported by Meena et al., (2017).The lowest CEC in treatment T₂ (Non-FYM) might be due to the acidifying effect of fertilizers resulting in reduced pH values in the 100% N-treated plots. While in sub-surface soil the highest CEC was obtained in T₃, FYM treated plot followed by T₁ and T₈. The lowest CEC was obtained in T₂ treated plot. CEC of all treatments in surface soil is higher than the subsurface soil. This may be due to organic matter content in surface soil is more than subsurface soil and organic matter could increase the CEC of the surface soil.

Table 5. Effect of nutrient management practices on exchangeable Ca, Mg and CEC of surface soil

Surface Soil 0-15 cm			
Treatments	Exch. Calcium (meq100g ⁻¹)	Exch. Mg (meq100g ⁻¹)	CEC [cmol (p+)kg ⁻¹]
T ₁ Control	4.1	2.8	11.8
T ₂ 100 % N	2.5	0.7	6.7
T ₃ FYM	6.2	3.8	13.2
T ₄ 100 % NP	3.1	0.9	9.84
T ₅ 100 % NPK	3.4	1	11.2
T ₆ ½ (N+FYM) + P (A-X/2) + K (B-Y/2) (INM)	5.3	3.2	12.7
T ₇ 100 % NPK + Lime	8.1	1.6	11.6
T ₈ Lime +FYM+P(A-X)+K(B-Y)	7.9	3.9	12.8
T ₉ Lime + N	5.7	2.1	8.2
CD(P=0.05)	0.37	0.36	0.27
CV(%)	4.07	9.38	1.54

Table 6. Effect of nutrient management practices on exchangeable Ca, Mg and CEC of sub-surface soil

Subsurface soil 16-30 cm				
Treatments		Exch. Calcium (meq100g ⁻¹)	Exch. Mg (meq100g ⁻¹)	CEC [cmol (p+)kg ⁻¹]
T ₁	Control	3.8	1.1	10.9
T ₂	100 % N	2.1	0.13	6.08
T ₃	FYM	5.9	3.6	11.6
T ₄	100 % NP	2.9	0.4	8.73
T ₅	100 % NPK	3	0.7	10.3
T ₆	½ (N+FYM) + P (A-X/2) + K (B-Y/2) (INM)	4.8	2.5	11.6
T ₇	100 % NPK + Lime	7.7	1.3	10.2
T ₈	Lime+FYM+P(A-X)+K(B-Y)	7.6	2.7	10.9
T ₉	Lime + N	5.4	1.9	7.83
CD(P=0.05)		0.46	0.42	0.43
CV(%)		5.51	5.11	2.72

3.3 Correlation of Exchangeable K, Ca, Mg, CEC with Various Soil Parameters

3.3.1 Correlation studies of exchangeable K, Ca, Mg, CEC with various soil parameters (0-15 cm depth)

The correlation coefficient (r) values worked out among the various soil parameters at two different depth of soil (0-15 cm and 16-30 cm) in the Maize Wheat cropping system.

From the table (Table 7) it was evident that the pH was found to be highly significant and positively correlated with Exchangeable Calcium (r=0.863**). Soil colloids acquire their charges either by dissociation of H⁺ from or onto the active functional group [19] and as the pH of the soil increases, the H⁺ ions on the functional groups of organic matter dissociate, resulting in the creation of negative charges on the organic matter. The negative charges created therefore have affinity for K⁺ ions, hence the positive correlation between organic carbon and the K forms observed. Amoakwah and Frimpong [20]. The parameter Electrical conductivity was observed to be non-significant and positively correlated with all the parameters under study. Organic Carbon was significant and positively correlated with Exchangeable Magnesium, Available K, Water soluble K, and Exchangeable K (r=0.780**, 0.771**, 0.748**, & 0.773** respectively). All the K forms positively correlated (P < 0.05) with organic carbon. This observation contradicts the findings of Acquaye (1973) who found no significant correlation of K with percentage organic carbon. Water Soluble and exchangeable K forms positively correlated with

organic matter, with correlation coefficients (r) of 0.748* and 0.773* for water soluble K and exchangeable K respectively. This observation may be attributed to the fact that, tropical soils are often characterized by a variable charge systems (i.e., pH dependent charges) [19] indicative that the creation of charges on tropical soil colloids depends on soil pH. Thus, the charge generated could either be positive or negative depending on the pH of the soil. Soil colloids acquire their charges either by dissociation of H⁺ from or onto the active functional group [19] and as the pH of the soil increases, the H⁺ ions on the functional groups of organic matter dissociate, resulting in the creation of negative charges on the organic matter. The negative charges created therefore have affinity for K⁺ ions, hence the positive correlation between organic carbon and the K forms observed and similar findings reported by Amoakwah and Frimpong [20]. Exchangeable Calcium was highly significant and positively correlated with pH (r=0.863**). Exchangeable Magnesium was significantly and positively correlated with Organic Carbon (r=0.780*). Cation exchange capacity (CEC) was found to be non-significantly and positively correlated with all the other parameters under study. Percentage K saturation values found in the soils also were high, but the overall low CEC values measured in the soil suggest that the soils are not adequately supplied with available K. It can therefore be concluded that potash fertilization would be required for sustainable crop production in the soils studied. Amoakwah and Frimpong [20]. Available K was found to be highly significant and positively correlated with Water soluble K and Exchangeable K (r=0.997** & 0.997**, respectively). In agreement with Samadi et al. [21]

the study showed significantly ($P < 0.05$) positive correlation ($r = 0.977^*$) between water soluble K and exchangeable, and between water soluble K and available K ($r = 0.977^*$, $P < 0.05$). This observation was not unexpected because exchangeable potassium is usually released into the soil solution from the exchange complex when plants deplete the soluble potassium, indicative that the size of the exchangeable potassium pool will determine the effectiveness of K re-supply, as well as the concentration of K in the soil solution. Similar to findings made by Ghiri et al. [22], also made similar observations to confirm the positive correlation previously found between exchangeable and available forms of potassium by Kasker et al. [23] and Kozak et al. [24]. Water soluble K was highly significant and positively correlated with available K, exchangeable K ($r=0.997^{**}$ & 0.997^{**} , respectively). It was significant and positively correlated with organic carbon ($r=0.748^*$). In agreement with Samadi et al. [21] the study showed “significantly ($P < 0.05$) positive correlation ($r = 0.997^*$) between water soluble K and exchangeable, and between water soluble K and available K ($r = 0.997^*$, $P < 0.05$). This observation was not unexpected because exchangeable potassium is usually released into the soil solution from the exchange complex when plants deplete the soluble potassium, indicative that the size of the exchangeable potassium pool will determine the effectiveness of K re-supply, as well as the concentration of K in the soil solution”. Similar to findings made by Ghiri et al. [22], exchangeable K concentrations in the soils positively correlated with available K ($r = 0.997^*$, $P < 0.05$) also made similar observations to confirm the positive correlation previously found between exchangeable and available forms of potassium by Kasker et al. [23] and Kozak et al. [24] and Amoakwah and Frimpong [20]. Exchangeable potassium was highly significant with Available K ($r=0.997^{**}$) and with Water soluble K ($r=0.997^{**}$). Whereas it was significantly and positively correlated with organic carbon ($r=0.773^*$). Exchangeable-K showed positive relationship with Organic Carbon ($r =0.773^*$) which is very much expected because of the fact that higher content of Organic Carbon in the soil leads to higher CEC resulting in higher adsorption of the cations including K [25]. All the K forms positively correlated ($P < 0.05$) with organic carbon. This observation contradicts the no significant correlation of K with percentage organic carbon. Soluble and exchangeable K forms positively correlated with organic matter, with correlation coefficients (r) of 0.567 and

0.701 for water soluble K and exchangeable K respectively. This observation may be attributed to the fact that, tropical soils are often characterized by a variable charge systems (i.e., pH dependent charges) [19], indicative that the creation of charges on tropical soil colloids depends on soil pH. “These results have indicated the existence of dynamic equilibrium between forms of K in soils of western plain of Rajasthan” (Dixit et al., 1993) [26,14].

3.3.2 Correlation studies of exchangeable Ca, Mg, CEC and potassium with various soil parameters (16-30 cm depth)

From the (Table 8) it was evident that the pH was found to be highly significant and positively correlated with Exchangeable Calcium ($r=0.879^{**}$), and was significant and positively correlated with Exchangeable Magnesium ($r=0.669^*$). The parameter Electrical conductivity was observed to be non-significant and positively correlated with all the parameters under study. Organic carbon was significant and positively correlated with Exchangeable Magnesium, Available K, Water soluble K, and Exchangeable K ($r=0.729^*$, 0.751^* , 0.751^* , & 0.753^*). Exchangeable Calcium was highly significant and positively correlated with pH ($r=0.879^{**}$). It was significant and positively correlated with Exchangeable Magnesium ($r=0.685^*$). Exchangeable Magnesium was significantly and positively correlated with pH, Organic carbon and Exchangeable Calcium ($r=0.669^*$, 0.729^* & 0.685^*) respectively. Cation exchange capacity (CEC) was found to be non-significantly and positively correlated with all the other parameters under study. Available K was found to be highly significant and positively correlated with Water soluble K and Exchangeable K ($r=0.999^{**}$ & 0.997^{**} respectively). It was significant and positively correlated with Organic carbon ($r=0.754^*$) respectively. Water soluble K was highly significant and positively correlated with Available K and with Exchangeable K ($r=0.999^{**}$ & 0.999^{**}) respectively. And it was significant and positively correlated with Organic carbon ($r=0.754^*$ & 0.796^*) respectively. Exchangeable K in the surface soil depth correlated with exchangeable K in the 16-30 cm soil layer. Significant and positive correlations between water-soluble K and exchangeable K forms indicate the faster rate of equilibrium between these forms. Similar interrelationships have been reported by Singh et al., (1993), Dixit et al., (1993), and Dan et al. [27]. Significant correlations between different forms indicate the

Table 7. Correlation coefficient of different nutrients in surface soil under Maize-wheat cropping system

	pH	EC (dSm ⁻¹)	O.C.(g kg ⁻¹)	Exch. Calcium (meq 100g ⁻¹)	Exch.Mg (meq 100g ⁻¹)	CEC cmol (p ⁺) kg ⁻¹	Available K (kg ha ⁻¹)	Water soluble K(kg ha ⁻¹)	Exchangeable K (kg ha ⁻¹)
pH	1								
EC(dSm ⁻¹)	-0.357	1							
O.C. (g kg ⁻¹)	0.292	0.280	1						
Exch. Calcium (meq 100g ⁻¹)	0.863**	0.045	0.448	1					
Exch. Mg (meq 100g ⁻¹)	0.644	-0.136	0.780*	0.628	1				
CEC cmol(p ⁺) kg ⁻¹	0.034	-0.314	0.609	0.066	0.614	1			
Available K (kg ha ⁻¹)	0.180	0.437	0.771*	0.301	0.462	0.167	1		
Water soluble K(kg ha ⁻¹)	0.141	0.439	0.748*	0.256	0.424	0.157	0.997**	1	
Exchangeable K (kg ha ⁻¹)	0.185	0.436	0.773*	0.307	0.466	0.168	0.997**	0.997**	1

**Significant at 1%level of significance

*Significant at 5%level of significance

Table 8. Correlation coefficient of different nutrients in sub-surface soil under Maize-wheat cropping system

	pH	EC (dSm ⁻¹)	O.C. (g kg ⁻¹)	Exch. Calcium (meq 100g ⁻¹)	Exch.Mg (meq 100g ⁻¹)	CEC cmol(p ⁺) kg ⁻¹	Available K (kg ha ⁻¹)	Water soluble K(kg ha ⁻¹)	Exchangeable K (kg ha ⁻¹)
pH	1								
EC(dSm ⁻¹)	-0.172	1							
O.C. (g kg ⁻¹)	0.263	0.297	1						
Exch. Calcium (meq 100g ⁻¹)	0.879**	0.240	0.413	1					
Exch.Mg ((meq 100g ⁻¹)	0.669*	0.126	0.729*	0.685*	1				
CEC cmol(p ⁺) kg ⁻¹	0.015	-0.270	0.608	0.005	0.463	1			
Available K Kg ha ⁻¹	0.168	0.369	0.754*	0.298	0.348	0.172	1		
Water soluble K(Kg ha ⁻¹)	0.166	0.384	0.751*	0.296	0.351	0.163	0.999**	1	
Exchangeable K (Kg ha ⁻¹)	0.168	0.367	0.753*	0.297	0.346	0.173	0.997**	0.999**	1

**Significant at 1%level of significance

*Significant at 5%levelofsignificance

existence of dynamic equilibrium among different K fractions. Thus, it can be postulated that each form of K influences another directly or indirectly. Hence, all forms are important in one way or other for K availability in soil (Girija and Badrinath, 1996). Similar finding was reported by Lakaria et al. [6]. Exchangeable K was highly significant with Available K and with Water soluble ($r=0.997^{**}$ & 0.999^{**} , respectively). Whereas it was significantly and positively correlated with Organic Carbon ($r=0.745^{*}$) respectively [28-30].

4. CONCLUSIONS

The results indicated that exchangeable K was found to be significantly highest in T_8 than rest of the treatments followed by T_7 and T_5 while lowest was in T_4 treatment in both depths respectively. The exchangeable Ca^{2+} was found to be highest in treatment T_7 followed by T_8 and T_3 than rest of the other treated plot while lowest amount was found in T_2 in both the depths. The trend of exchangeable magnesium was found to be significantly different in 0-15 cm and 16-30 cm soil depths as the highest amount was observed in treatments such as T_8 followed by T_3 and T_6 in surface depths (0-15 cm) while in sub-surface depths (15-30 cm) the highest amount was found in treatments T_3 followed by T_8 and T_6 while the lowest amount was found in T_2 treated plot in both depths. Cation exchangeable capacity (CEC) is one the vital soil parameters concerning to soil fertility measured in terms of $cmol(p+)$ kg^{-1} of soil and it is significantly varied in surface soils as well as in sub-surface soil depths. The CEC was significantly higher in FYM treated plot as compared to Non-FYM treated plot. Therefore, the highest CEC was recorded in T_3 treated plot than rest of the other treatment while the lowest one was recorded in T_2 plot in surface soils as well as sub-surface soils. The correlation coefficient (r) values worked out among the various soil parameters at these depth of soil where pH was significantly and positively correlated with exchangeable Ca^{2+} and exchangeable Mg^{2+} . Similarly organic carbon shows positive correlation with exchangeable Mg^{2+} and exchangeable K^+ . Whereas EC and CEC shows non-significant but positive correlation with the parameters taken under study. Hence, the result of the present study will be highly useful for improving the soil fertility by using suitable and judicious combination of nutrient management with various cropping system side by side and hence provides a scientific scope for further research work.

ACKNOWLEDGEMENTS

The authors express their sincere gratitude and thanks to Dr. Dharendra Kumar Shahi, Head of the Department of Soil Science and Dean of Birsa Agricultural University, Kanke, Ranchi, Dr. Prabakar Mahapatra, Junior Scientist-cum-Assistant Professor and all the supportive member of the department for their valuable and constructive suggestions during the planning and development of the research work and its implementation. The necessary facilities provided by the department of soil science and collaboration with other department during the itinerary of research work is highly acknowledge.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Davari MR, Mirzakhani M. Integrated nutrient management towards sustainable production of oilseeds and pulses. International Journal of Agriculture and Crop Science. 2009;1:24-32.
2. Kannan RL, Dhivya M, Abinaya D, Lekshmi Krishna R and Krishnakumar S. Effect of integrated nutrient management on soil fertility and productivity in maize. Bulletin of Environment, Pharmacology and Life Sciences. 2013;2:61-67.
3. Singh B, Singh Y. Concepts in nutrient management. In: Recent Advances in Agronomy. Indian Soc. Agron., New Delhi. 2002;92-109.
4. Gawai PP, Pawar VP. Integrated nutrient management in sorghum (*Sorghum bicolor*)-chickpea (*Cicer arietinum*) cropping sequence under irrigated conditions. Indian J Agron. 2006;51: 17-20.
5. Baruah TC, Barthakur HP. A textbook of soil analysis. Vikas Publishing House Pvt. Ltd; 1999.
6. Lakaria BL, Behera SK, Singh D. Different forms of potassium and their contributions towards potassium uptake under long-term maize (*Zea mays* L.)-Wheat (*Triticum aestivum* L.)- Cowpea (*Vigna unguiculata* L.) rotation on an inceptisol. Communication of Soil Science and Plant Analysis. 2012;43:936-947.

7. Mazumdar SP, Kundu DK, Ghosh D, Saha AR, Majumdar B, Ghorai AK. Effect of long-term application of inorganic fertilizers and organic manure on yield, potassium uptake and distribution of potassium fractions in the new gangetic alluvial soil under jute-rice-wheat cropping system. *International Journal of Agriculture and Food Science Technology*. 2014;5(4):297-306. ISSN 2249-3050
8. Sharma U, Paliyal SS. Forms of soil potassium as influenced by long term application of chemical fertilizers and organics in rainfed maize-wheat cropping system. *Journal of Krishi Vigyan*. 2015;3(2):48-53.
9. Jatav MK, Sud KC, Trehan SP. Effect of organic and inorganic sources of phosphorus and potassium on their different fractions under potato-radish cropping sequence in a Brown Hill Soil. *Journal of the Indian Society of Soil Science*. 2010;58:388-393.
10. Kumari PD. Effect of crop residue incorporation in hybrid maize on k fraction in acid soil of Ranchi, M.Sc. Thesis. Submitted to SSAC, BAU, Ranchi; 2017.
11. Bhattacharyya R, Prakash V, Kundu S, Ghosh BN, Srivastava AK, Gupta H. Potassium balance as influenced by farmyard manure application under continuous soybean-wheat cropping system in a Typic Haplaquept. *Geoderma*. 2006;137:155-160.
12. Sharma A, Jalali VK, Vivek M, Arya, Rai P. Distribution of various forms of potassium in soils representing intermediate zone of Jammu region. *Journal of the Indian Society of Soil Science*. 2009;57(2):205-207.
13. Sharma KN, Singh H, Bhandari AL. Influence of long-term fertilizer on K adsorption kinetics parameters, *Journal of Potassium Research*. 1994;10:368-379.
14. Prasad J. Forms of potassium in shallow soils of different origin and land uses in Nagpur district of Maharashtra. *Journal of the Indian Society of Soil Science*. 2010;58(3):327-330.
15. Babu MVS, Reddy CM, Subramanyam A, Balaguravaiah D. Effect of integrated use of organic and inorganic fertilizers on soil properties and yield of sugarcane. *Journal of the Indian Society of Soil Science*. 2007;55:161-166.
16. Brady NC, Weil RR. The nature and properties of soils. Thirteenth edition. Pearson Education (Singapore) Pte. Ltd., Indian Branch. New Delhi; 2002.
17. Verma S. Long term effect of nutrient management practices on soil quality parameters under maize-wheat cropping system. M.Sc. thesis. Submitted to SSAC, BAU, Ranchi; 2017.
18. Prasad J, Karmakar S, Kumar R, Mishra B. Influence of integrated nutrient management on yield and soil properties in maize-wheat cropping system in an Alfisol of Jharkhand. *Journal of the Indian Society of Soil Science*. 2010;58(2):200-204.
19. Van Ranst E. Properties and management of soils of the tropics. Chapter. 2006;2:31-32.
20. Amoakwah E, Frimpong KA. Relationship between potassium forms and selected physico-chemical properties of some Ghanaian soils along a toposequence *ARPN Journal of Engineering and Applied Sciences*. 2013;8(7).
21. Samadi A, Rezapour S, Jafarzadeh AA, Oustan S. Impact of clay mineralogy and landscape on potassium forms in calcareous soils, urmia region. *Journal Agricultural Science and Technology*. 2010;12:495-507.
22. Ghiri MN, Ali Abtahi A, Owliaie H, Soheila SH, Koohkan H. Arid land research and management. *Arid Land Research and Management*. 2011;25:313-327.
23. Kaskar DR, Salvi VG, Mayekar BS, Dabke DJ. Forms of potassium, their interrelationship and relationships with other soil properties of inceptisols of west coast of Maharashtra. *Journal of Potassium Research*. 2001;17:23-27.
24. Kozak M, Stêpień M, Joarder AH. Relationships between available and exchangeable potassium content and other soil properties. *Polish Journal of Soil Science*. 2005;38(2):1-8.
25. Shankhayan SD, Sharma HR, Bhardwaj SK. Response of wheat to potassium application in soils of una and paonta valleys of Himachal Pradesh. *Journal of Potassium Research*. 1996;12(3):290-297.
26. Chand S, Swami BN. Different forms of potassium in some important soil association of Bharatpur district of Rajasthan. *Journal of potassium Research*. 2000;16:59-61.
27. Dan S, Khurana HS, Thind SS. Forms of soil K and their contribution to plant k in Typic Ustipsament loamy sand soil under

- potato. *Journal of Potassium Research*. 2004;20:28-33.
28. Gomez KA, Gomez AA. *Statistical procedures for agricultural research*. John Wiley and Sons, New York; 1984.
29. Sharma SP, Singh MV, Subehia SK, Jain PK, Kaushal V, Verma TS. Long term effect of fertilizer, manure and lime application on the changes in soil quality, crop productivity and sustainability of maize-wheat system in alfisol of North Himalaya. *Research Bulletin No.2. AICRP on Long Term Fertilizer Experiments, IISS, Bhopal (M.P) and Department of Soils, CSK HPKV, Palampur, H.P.* 2005;1-88.
30. Singh K, Aulakh MS. Nitrogen immobilization-mineralization in a sandy loam soil (Typic Ustochrept) amended with crop residues at varying moisture regimes. *Journal of the Indian Society of Soil Science*. 2001;49:198-202.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/116989>