



Response of Nano Dap on Yield and Nutrient Content of Wheat Grown in Loamy Sand Soil

Parmar G S^{a++*}, Viradiya M B^{b#} and Patel J A^{at}

^a Department of Soil Science and Agricultural Chemistry, Anand Agricultural University, Anand, Gujarat-388 110, India

^b Seth M. C. Polytechnic in Agriculture, Anand Agricultural University, Anand, Gujarat-388 110, India

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i84866>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120847>

Original Research Article

Received: 25/05/2024

Accepted: 27/07/2024

Published: 30/07/2024

ABSTRACT

A field experiment was conducted at Agronomy Farm, B. A. College of Agriculture, Anand Agricultural University, Anand to study the "Response of Nano DAP on Yield and Nutrient Content of Wheat Grown in Loamy Sand Soil" in *rabi* season during the year 2023-24. The experiment consisted of ten treatments and was arranged in a randomized block design with three replications. The results proved that the combined application of conventional and nano DAP had a positive impact on most of the growth parameters. The application of 100% RDF N and P (T₁), (75% RDF N and P + foliar spray with nano DAP @ 4 ml/L at 25 DAS) (T₆) and 50% RDF N and P + Seed treatment with nano DAP @ 5 ml/kg seed + First foliar spray with nano DAP @ 4 ml/L at 25

⁺⁺ M.Sc. Scholar;

[#] Principal and Associate Professor;

[†] PhD. Scholar;

*Corresponding author: E-mail: parmargaurang889@gmail.com;

Cite as: G S, Parmar, Viradiya M B, and Patel J A. 2024. "Response of Nano Dap on Yield and Nutrient Content of Wheat Grown in Loamy Sand Soil". *International Journal of Plant & Soil Science* 36 (8):374-83. <https://doi.org/10.9734/ijpss/2024/v36i84866>.

DAS(T₁₀) obtained significantly higher plant height, root length, root biomass, effective numbers of tillers, spike length, test weight, grain yield, straw yield. Similarly, nutrient content and uptake of N and P by grain and straw under T₂ and T₆ were significantly higher than rest of the treatments. Therefore, we may suggest for 75% RDF N and P + foliar spray with nano DAP @ 4 ml/L at 25 DAS and 50% RDF N and P + Seed treatment with nano DAP @ 5 ml/kg seed + First foliar spray with nano DAP @ 4 ml/L at 25 DAS for better realization of applied nutrients in terms yield responses.

Keywords: Nano DAP; foliar application.

1. INTRODUCTION

Wheat scientifically known as *Triticum aestivum* L., belongs to the Poaceae family and serves as a fundamental crop globally, holding particular significance in India as a primary winter crop. Wheat cultivation covered approximately 31.40 million hectares, resulting in a total production of around 110.55 million tonnes with average wheat productivity was recorded at 3248 kg/ha in 2022-23 [1]. Nano fertilizers are synthesized or modified forms of traditional fertilizers, bulk materials or extracts from various vegetative or reproductive parts of plants. Nano fertilizers play a crucial role in enhancing nutrient uptake and nutrient use efficiency, reducing losses through leaching and gaseous emissions and minimizing the risk of nutrient toxicity. This contributes to ensuring food security achieved through higher productivity and economic returns by practicing sustainable farming practices [2].

Nano DAP is a highly efficient source of nitrogen (N) and phosphorus (P₂O₅) for all types of crops, effectively addressing deficiencies in these nutrients in standing crops. The nano DAP formulation contains 8% nitrogen and 16% phosphorus (P₂O₅) by weight/volume. Its liquid form benefits from a high surface-area-to-volume ratio, as its particle size is under 100 nanometers (nm). The nitrogen and phosphorus nano clusters in nano DAP are functionalized with biopolymers and other excipients, enhancing their spreadability and assimilation within the plant system. This results in higher seed vigor, increased chlorophyll and photosynthetic efficiency, improved crop quality, and higher yields [3].

2. MATERIALS AND METHODS

The experiment was carried at Agronomy Farm, B. A. College of Agriculture, Anand Agricultural University, Anand, India during *rabi* season of the year 2023-24 to examine the response of nano dap on yield and nutrient content of wheat. The experimental site had a smooth terrain with slight

slopes and effective drainage. The soil type known as "Goradu" soil, is a true depiction of soils found in the middle Gujarat region. It is characterized by a loamy sand texture, good moisture retention and belongs to the Inceptisols soil order. wheat (variety-GW 496) was used as test crop in the experiment. The trial was laid down in a randomized block design (RBD) corresponding to 10 treatments and three replications. The details of treatments are as follows, T₁ (Absolute control), T₂ (100% RDF N and P), T₃ (75% RDF N and P), T₄ (50% RDF N and P), T₅ (T₃ + Foliar spray with Nano DAP @ 2 ml/L at 25 DAS), T₆ (T₃ + Foliar spray with Nano DAP @ 4 ml/L at 25 DAS), T₇ (T₄ + Foliar spray with Nano DAP @ 2 ml/L at 25 DAS), T₈ (T₄ + Foliar spray with Nano DAP @ 4 ml/L at 25 DAS), T₉ (T₄ + Seed treatment with Nano DAP @ 2.5 ml/kg seed + First foliar spray with Nano DAP @ 2 ml/L 25 DAS) and T₁₀ (T₄ + Seed treatment with Nano DAP @ 5 ml/kg seed + First foliar spray with Nano DAP @ 4 ml/L 25 DAS). Recommended basal dose of phosphorus (60 kg/ha) and nitrogen (120 kg/ha) was applied as per treatment *i.e.*, 100%, 75% and 50% RDF. Half dose of 100%, 75% and 50% RDF was supplied through diammonium phosphate (DAP) and urea before sowing of seeds as basal application and the remaining half dose of nitrogen through urea was applied as top dressing at CRI stage. while, nano DAP was given at 25 DAS through foliar application, as per the treatments.

The Statistical analysis of various parameters such as growth, yield, and quality were conducted using computer-based methods. The statistical assessment followed the Randomized Block Design and conducted at the Department of Agricultural Statistics, B. A. College of Agriculture, AAU, Anand in accordance with the procedures specified by Cochran and Cox (1967). The variances originating from different sources in the analysis of variance (ANOVA) were examined using the "F-test" and compared with the value of F at 5% level of significance. The calculations for standard error of the mean

(S. Em±), Critical Differences (CD), and Coefficient of Variation (CV %) were also determined.

3. RESULTS AND DISCUSSION

3.1 Plant Height

The applications of both conventional DAP and its combination with foliar spray of nano DAP had significant results on the plant height at 30 DAS and harvest. The treatment T₂ (100% RDF N and P) recorded significantly higher plant height (44.09 cm) at 30 DAS and it was remained at par with treatment T₆ (75% RDF N and P + foliar spray with nano DAP @ 4 ml/L at 25 DAS). While significantly higher plant height (105.51 cm) at harvest observed with T₂ (100% RDF N and P) and it was remained at par with treatments T₅, T₆ and T₁₀. Significantly lower plant height of 34.12 cm and 80.75 cm were recorded in treatment T₁ (Absolute control) at 30 DAS and harvest, respectively. The increase in plant height at 30 DAS and at harvest of wheat with the application of nano DAP might be due to the smaller size of nano DAP particles compared to regular fertilizer, allowing for better absorption by plants. This provides a readily available source of phosphorus (P) and nitrogen (N), which are essential for stem growth. This result is consistent with the findings of Sahithi et al. [4] Chinnappa et al. [5] and Prakash et al. [6].

3.2 Root Length and Root Biomass

The treatment T₂ (100% RDF N and P) recorded significantly higher root length (11.58 cm) at 30-35 DAS and it was remained at par with treatment T₅, T₆ and T₁₀. In contrast, the lower root length (8.71 cm) was recorded with treatment T₁ (Absolute control). The treatment T₂ (100% RDF N and P) recorded significantly higher root biomass (4.19 g/plant) at 30-35 DAS and it was remained at par with treatment T₅ and T₆. While the lower root biomass (2.41 g/plant) was recorded with treatment T₁ (Absolute control). These results were in close agreement with those found by Abdel-Aziz et al. [7] Amira et al. [8] and Maloth et al. [9].

3.3 Number of Effective Tillers/m Row Length

Number of effective tillers per meter row length (108.25) at harvest was found significantly higher with treatment T₂ (100% RDF N and P) as compared to other treatments and was found to

be at par with the treatments T₅, T₆ and T₁₀. In contrast, the lower number of effective tillers (89.49) per meter row length was recorded with treatment T₁ (Absolute control). The increase in the total number of tillers and effective tillers with nano DAP might be attributed to its smaller particle size enhances nutrient uptake, facilitating more efficient absorption of essential nutrients such as phosphorus (P) and nitrogen (N). This efficient uptake supports better root development, providing a robust foundation for multiple tillers, and stimulates growth hormones that encourage tiller formation. Additionally, Nano DAP allows for more uniform nutrient distribution and enhances the plant's stress tolerance. Collectively, these factors lead to a significant increase in the number of effective tillers per plant. These results are in close accordance with those reported by Meena et al. [10] and Sahu et al. [11].

3.4 Spike Length

Significantly higher spike length (10.34 cm) with treatment T₂ (100% RDF N and P) compared to other treatments. This treatment was found to be statistically at par with treatments T₅, T₆, T₈ and T₁₀. The lower spike length (8.64 cm) was observed in treatment T₁ (Absolute control). These results closely align with those reported by Gomaa et al. [12].

3.5 Test Weight

The treatment T₂ (100% RDF N and P) recorded significantly higher test weight of wheat grain (42.52 g) than other treatments but was found statistically at par with the treatments T₃, T₅, T₆, T₈, T₉ and T₁₀. While treatment T₁ (Absolute control) recorded the lower test weight (36.29 g) as compared to other treatments.

3.6 Grain Yield

The treatment T₂ (100% RDF N and P) recorded significantly higher grain yield of wheat (6315kg/ha) than other treatments and it was at par with treatment T₅, T₆ and T₁₀. In contrast lower grain yield of wheat (4425 kg/ha) was recorded with the treatment T₁ (Absolute control). The rise in wheat grain yield is mainly attributed to the improved growth of plant parts and metabolic activities like photosynthesis, resulting in increased accumulation and translocation of photosynthates to the plant's economic parts. Nano fertilizers raised broad bean production and yield components as a

result of increased growth hormone activity and improved metabolic processes, which tend to increase flowering and grain formation (El-Azizy et al., 2021). Similar results were also reported by Rajput et al. [13] and Sahu et al. [11].

3.7 Straw Yield

The treatment T₂ (100% RDF N and P) recorded significantly higher straw yield of wheat (8114 kg/ha) than other treatments and it was at par with treatment T₅, T₆ and T₁₀. In contrast lower straw yield of wheat (5503 kg/ha) was recorded with the treatment T₁ (Absolute control). The increased grain and straw yield of wheat after the application of nano fertilizers may be attributed to increased growth hormone levels, enhanced metabolic processes and improved photosynthetic activities due to the application of nano fertilizers [10]. Similar results were found by Reddy et al. [14] and Prakash et al. [6].

3.8 Protein Content (%)

The treatment T₂ (100% RDF N and P) recorded significantly higher protein content of wheat grain (11.94%) than other treatments but was found statistically at par with the treatments T₅, T₆, T₈ and T₁₀. Treatment T₁ (Absolute control) recorded the lower protein content of wheat grain (10.54%) as compared to other treatments. Nano DAP increases the protein content of wheat by enhancing nitrogen and phosphorus availability, which are crucial for protein synthesis and energy transfer. Its smaller particle size allows for more efficient nutrient absorption, boosting photosynthesis and overall plant health. This leads to better protein synthesis and accumulation in the wheat grains. These results are in conformity to those reported by Goma et al. [12] and Dhansil et al. [15].

3.9 Nitrogen Content

The treatment T₂ (100% RDF N and P) recorded significantly higher N content of wheat grain (1.91%) than other treatments but was found statistically at par with the treatments T₅, T₆ and T₁₀. Treatment T₁ (Absolute control) recorded the lower N content of wheat grain (1.69%) as compared to other treatments while treatment T₂ (100% RDF N and P) recorded significantly higher N content of wheat straw (0.76%) than other treatments but was found statistically at par with the treatments T₅, T₆ and T₁₀. In

contrast treatment T₁ (Absolute control) recorded lower N content in wheat straw (0.66%) compared to other treatments.

3.10 Phosphorus Content

The treatment T₂ (100% RDF N and P) recorded significantly higher P content of wheat grain (0.390%) than other treatments but was found statistically at par with the treatments T₅, T₆, T₈, T₉ and T₁₀. Treatment T₁ (Absolute control) recorded the lower P content of wheat grain (0.327%) as compared to other treatments while treatment T₂ (100% RDF N and P) recorded significantly maximum P content of wheat straw (0.187%) than other treatments but was found statistically at par with the treatments T₃, T₅, T₆, T₈, T₉ and T₁₀. Treatment T₁ (Absolute control) recorded lower P content in wheat grain (0.163%) compared to other treatments [16,17].

3.11 Nitrogen Uptake by Grain and Straw

Nitrogen uptake by grain and straw of wheat was found to be significant under different nano DAP treatments. Significantly higher nitrogen uptake by grain and straw (122.49 kg/ha and 70.92 kg/ha respectively) was recorded with the treatment T₂ (100% RDF N and P) and was found to be at par with the treatments T₅, T₆ and T₁₀. Treatment T₁ (Absolute control) recorded the lower N uptake of wheat grain and straw (76.48 kg/ha and 45.46 kg/ha respectively) as compared to other treatments.

3.12 Phosphorus Uptake by Grain and Straw

Phosphorus uptake by grain and straw of wheat was found to be significant under different nano DAP treatments. Significantly higher phosphorus uptake by grain and straw (25.01 kg/ha and 17.69 kg/ha respectively) was recorded with the treatment T₂ (100% RDF N and P) and was found to be at par with the treatments T₅, T₆ and T₁₀. Treatment T₁ (Absolute control) recorded the lower P uptake of wheat grain and straw (14.76 kg/ha and 11.10 kg/ha respectively) as compared to other treatments.

3.13 Available N and P Content in Soil

The treatment T₂ (100% RDF N and P) recorded the higher available N (211 kg/ha) compared to other treatments but was statistically at par with treatments T₃, T₅ and T₆. whereas treatment T₁ (Absolute control) recorded significantly lower

Table 1. Effect of different treatments on plant height, root length, root biomass and no. of effective tillers

Tr No.	Treatments	Plant height(cm)		Root length(cm)	Root biomass (g/plant)	No. of effective tillers/m row length
		30 DAS	Harvest			
T ₁	Absolute Control	34.12	80.75	8.71	2.41	89.49
T ₂	100% RDF N and P	44.09	105.51	11.58	4.19	108.25
T ₃	75% RDF N and P	36.37	84.38	9.36	2.75	94.07
T ₄	50% RDF N and P	35.15	82.85	9.63	2.49	92.80
T ₅	T ₃ + Foliar spray with Nano DAP@ 2 ml/L at 25 DAS	38.06	98.31	11.03	3.95	104.69
T ₆	T ₃ + Foliar spray with Nano DAP@ 4 ml/L at 25 DAS	39.17	98.80	11.21	3.97	106.97
T ₇	T ₄ + Foliar spray with Nano DAP@ 2 ml/L at 25 DAS	35.30	82.91	9.33	2.73	93.87
T ₈	T ₄ + Foliar spray with Nano DAP@ 4 ml/L at 25 DAS	37.37	87.86	10.68	3.57	95.28
T ₉	T ₄ + Seed treatment with Nano DAP @ 2.5 ml/kg seed + First foliar spray with Nano DAP@ 2 ml/L 25 DAS.	37.11	86.05	9.80	3.36	95.26
T ₁₀	T ₄ + Seed treatment with Nano DAP @ 5 ml/kg seed + First foliar spray with Nano DAP@ 4 ml/L 25 DAS	37.42	96.54	10.85	3.57	104.03
	S. Em. ±	1.72	4.31	0.43	0.14	4.20
	C.D. at 5%	5.10	12.82	1.29	0.42	12.47
	C.V. (%)	7.95	8.27	7.34	7.39	7.38

Table 2. Effect of different treatments on spike length, grain yield, straw yield, test weight and protein content

Tr No.	Treatments	Spike length (cm)	Grain yield (kg/ha)	Straw yield (kg/ha)	Test weight (g)	Protein content (%)
T ₁	Absolute Control	8.64	4425	5503	36.29	10.54
T ₂	100% RDF N and P	10.34	6315	8114	42.85	11.94
T ₃	75% RDF N and P	9.09	5290	6471	38.03	11.02
T ₄	50% RDF N and P	8.74	4708	5783	36.33	10.75
T ₅	T ₃ + Foliar spray with Nano DAP@ 2 ml/L at 25 DAS	9.97	5810	7268	41.51	11.54
T ₆	T ₃ + Foliar spray with Nano DAP@ 4 ml/L at 25 DAS	10.18	6064	7869	42.12	11.79
T ₇	T ₄ + Foliar spray with Nano DAP@ 2 ml/L at 25 DAS	8.88	5221	6270	37.80	10.96
T ₈	T ₄ + Foliar spray with Nano DAP@ 4 ml/L at 25 DAS	9.22	5299	6536	37.84	11.19
T ₉	T ₄ + Seed treatment with Nano DAP @ 2.5 ml/kg seed + First foliar spray with Nano DAP@ 2 ml/L 25 DAS.	9.16	5291	6518	38.42	11.10
T ₁₀	T ₄ + Seed treatment with Nano DAP @ 5 ml/kg seed + First foliar spray with Nano DAP@ 4 ml/L 25 DAS	9.95	5664	6971	41.14	11.48
	S. Em. ±	0.40	333	524	1.46	0.27
	C.D. at 5%	1.18	988	1557	4.34	0.79
	C.V. (%)	7.32	10.65	13.49	6.45	4.09

Table 3. Effect of different treatments on nitrogen and phosphorus content in grain and straw as well as N and P uptake by grain and straw

Tr No.	Treatments	N content (%)		P content (%)		N uptake (kg/ha)		P uptake (kg/ha)	
		Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw
T ₁	Absolute Control	1.69	0.66	0.327	0.163	76.48	45.46	14.76	11.10
T ₂	100% RDF N and P	1.91	0.76	0.390	0.187	122.49	70.92	25.01	17.69
T ₃	75% RDF N and P	1.76	0.69	0.347	0.169	94.81	54.99	18.74	13.10
T ₄	50% RDF N and P	1.72	0.68	0.330	0.163	82.51	48.61	15.94	11.63
T ₅	T ₃ + Foliar spray with Nano DAP@ 2 ml/L at 25 DAS	1.85	0.74	0.377	0.180	109.36	64.02	22.27	15.48
T ₆	T ₃ + Foliar spray with Nano DAP@ 4 ml/L at 25 DAS	1.89	0.75	0.387	0.186	116.47	68.85	23.85	17.13
T ₇	T ₄ + Foliar spray with Nano DAP@ 2 ml/L at 25 DAS	1.75	0.67	0.340	0.167	93.20	53.82	18.21	12.65
T ₈	T ₄ + Foliar spray with Nano DAP@ 4 ml/L at 25 DAS	1.80	0.68	0.367	0.171	96.70	56.34	19.80	13.37
T ₉	T ₄ + Seed treatment with Nano DAP @ 2.5 ml/kg seed + First foliar spray with Nano DAP@ 2 ml/L 25 DAS	1.78	0.69	0.367	0.170	95.75	56.19	19.75	13.38
T ₁₀	T ₄ + Seed treatment with Nano DAP @ 5 ml/kg seed + First foliar spray with Nano DAP@ 4 ml/L 25 DAS	1.84	0.73	0.370	0.177	107.12	61.46	21.31	14.68
	S. Em. ±	0.04	0.02	0.01	0.01	6.63	3.55	1.43	0.99
	C.D. at 5%	0.11	0.05	0.03	0.02	19.69	10.54	4.25	2.93
	C.V. (%)	4.09	4.11	4.54	5.04	11.54	10.58	12.40	12.20

Table 4. Effect of different treatments on available nitrogen and phosphorus of soil after harvest of wheat

Tr No.	Treatments	Available nutrient status(kg/ha)	
		N	P ₂ O ₅
T ₁	Absolute Control	178.3	40.87
T ₂	100% RDF N and P	211.0	55.31
T ₃	75% RDF N and P	195.7	51.74
T ₄	50% RDF N and P	183.2	48.47
T ₅	T ₃ + Foliar spray with Nano DAP@ 2 ml/L at 25 DAS	200.0	52.23
T ₆	T ₃ + Foliar spray with Nano DAP@ 4 ml/L at 25 DAS	203.7	53.93
T ₇	T ₄ + Foliar spray with Nano DAP@ 2 ml/L at 25 DAS	184.7	46.31
T ₈	T ₄ + Foliar spray with Nano DAP@ 4 ml/L at 25 DAS	188.0	48.83
T ₉	T ₄ + Seed treatment with Nano DAP @ 2.5 ml/kg seed + First foliar spray with Nano DAP@ 2 ml/L 25 DAS	189.9	51.27
T ₁₀	T ₄ + Seed treatment with Nano DAP @ 5 ml/kg seed + First foliar spray with Nano DAP@ 4 ml/L 25 DAS	192.3	52.85
	S. Em. ±	5.32	1.39
	C.D. at 5%	15.81	4.12
	C.V. (%)	4.78	4.79

available nitrogen (178.3 kg/ha) as compared to other treatments. similarly, treatment T₂ (100% RDF N and P) recorded the higher available P₂O₅ (55.31 kg/ha) compared to other treatments but was statistically at par with treatments T₃, T₅ and T₆. In contrast treatment T₁ (Absolute control) recorded significantly lower available P₂O₅ (40.87 kg/ha) as compared to other treatments.

4. CONCLUSION

In light of the findings of the present investigation. It can be concluded that, the foliar application of nano DAP fertilizer in combination with conventional phosphatic fertilizer showed potential in reducing the soil application of phosphorus. It was observed that application of 100% RDF of N and P was superior. However, using 75% RDF N and P + foliar spray with nano DAP @ 4 ml/L at 25 DAS and 50% RDF N and P + Seed treatment with nano DAP @ 5 ml/kg seed + First foliar spray with nano DAP @ 4 ml/L at 25 DAS seems to be more effective for achieving higher growth and yield attributing characters of wheat.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous Centre for monitoring indian economy- commodities; 2023. Available:<https://commodities.cmie.com>
2. Ahmadian K, Jalilian J, Pirzad A. Nano-fertilizers improved drought tolerance in wheat under deficit irrigation. *Agricultural Water Management*. 2021;244:106544.
3. Anonymous. IFFCO Nano DAP fertilizer for sustainable farming; 2022. Available:<https://www.sagarika.com/en/nano-dap>
4. Sahithi B, Babu KK, Prasanth P, Lakshminarayana D, Gouthami P. Studies on effect of nano urea and nano dap alone and in combination with different levels of nitrogen and phosphorus on growth and yield of marigold (*Tagetes erecta* L.) Cv. superball. *The Pharma Innovation Journal*. 2023;12(11):901-906.
5. Chinnappa SA, Krishnamurthy D, Ajayakumar MY, Ramesha YM, Ravi S. Response of nano fertilizers on growth, yield and economics of kharif sorghum. *The Pharma Innovation Journal*. 2023;12(9):761-765.
6. Prakash AN, Siddaram R, Bellakki MA. Response of nano DAP on growth, yield and economics of Soybean (*Glycine max* L.). *The Pharma Innovation Journal*. 2023;12(12):1985-1989.
7. Abdel-Aziz H.M., Hasaneen, M.N., & Omer AM. Nano chitosan-NPK fertilizer enhances the growth and productivity of wheat plants grown in sandy soils. *Spanish Journal of Agricultural Research*. 2016;14(1):0902.
8. Amira Sh, Soliman M, Hassan FAE, Hanafy AH, Souad AEF. Effect of nano and molecular phosphorus fertilizers on growth and chemical composition of baobab (*Adansonia digitata* L.). *Journal of Plant Science*. 2016;11:52-60.
9. Maloth A, Thatikunta R, Parida BK, Naik DS, Varma N. Evaluation of nano-DAP on plant growth, enzymatic activity and yield in paddy (*Oryza sativa* L.). *International Journal of Environment and Climate Change*. 2024;14(1):890-897.
10. Meena M, Jat G, Meena RH, Sharma SK, Choudhary R, Yadav SK, Jain D, Choudhary RS, Chhippa BG, Jat H. Effect of phospho enriched compost and zinc on soil properties and productivity of blackgram (*Vigna mungo*). *Indian Journal of Agricultural Sciences*. 2021;91(9):1333-6.
11. Sahu D, Banwasi R, Shrivastava LK, Chowdhury T, Jatav G, Kumar H. Performance of Nano-DAP under Rice-wheat cropping system in Vertisol of Chhattisgarh, India. *International Journal of Environment and Climate Change*. 2023;13(9):2864-2875.

12. Gomaa MA, Radwan FI, Kandil EE, Al-Msari MAF. Response of some Egyptian and Iraqi wheat cultivars to mineral and nano fertilization. *Egyptian Academic Journal of Biological Sciences, H. Botany*. 2018;9(1):19-26.
13. Rajput JS, Thakur AK, Nag NK, Chandrakar T, Singh DP. Effect of nano fertilizer in relation to growth, yield and economics of little millet (*Panicum sumatrense* Roth) under rainfed conditions. *The Pharma Innovation Journal*. 2022;11(7):153-156.
14. Reddy BM, Elankavi S, Kumar MS, Sai MV, Vani BD. Effects of conventional and nano fertilizers on growth and yield of maize (*Zea mays* L.). *Bhartiya Krishi Anusandhan Patrika*. 2022;37(4):379-382.
15. Dhansil A, Zalawadia N, Prajapat BS, Yadav K. Effect of nano phosphatic fertilizer on nutrient content and uptake by pearl millet (*Pennisetum glaucum* L.) crop. *International Journal of Current Microbiology and Applied Science*. 2018;7(12):2327-2337.
16. Cochran WG, Cox GM. *Experimental designs*. North Carolina State University. Dept. of Statistics; 1948.
17. Singh MD. Nano fertilizer is a New Way to Increase Nutrient Use Efficiency in Crop Production. *International Journal of Agriculture Science*. 2017;9(7):3831-3833.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/120847>